


Vegetation Monitoring

August 2013

SAN JOAQUIN RIVER
RESTORATION PROGRAM



Vegetation Monitoring

In order to evaluate the establishment and development of riparian vegetation in response to Interim Flows, Reclamation's Technical Service Center in Denver, CO and Mid-Pacific Region in Sacramento, CA established monitoring transects in river reaches 1A through 5 and including the East Side and Mariposa Bypasses. In 2011, twenty permanent vegetation transects were established within river reaches 1A, 1B, 2A, 2B, 3, 4A, 4B2 (i.e. San Luis National Wildlife Refuge), and the East Side and Mariposa Bypasses. In 2012, two additional permanent vegetation transects were established within river reach 5. Due to the large project area (over 150 RM), it was feasible to locate and monitor two transects within each reach with the exception of the East Side Bypass, where four transects were placed, two of which were in the Merced NWR. Transects were placed in areas adjacent to the river channel within the active floodplain. These sites are subsequently subject to seasonal changes in water and nutrient input and scour and sediment deposition. These transects are not representative of vegetation types across entire reaches, but are illustrative of vegetation change over time resulting from Interim Flows. Monitoring of the transects for 2013 was completed in late June - no data analysis has been completed at this time.

Plant cover, composition, and overstory height and stem density were collected along each transect. The length of each transect was determined by the extent of the floodplain and varied from 35 to 100 meters (m). For herbaceous understory measurements, cover and species composition were measured either every 0.5 or 1 m along the transect depending on the length of the transect. The point-intercept method was used, which entailed recording the first "hit" for herbaceous plants by species and for woody species under 1 m tall. If a plant was not intercepted, then bare soil, litter, rock, or water were recorded. The location and extent of invasive weed species were documented when encountered.

The line-intercept method was used for measuring woody overstory cover. Overstory cover was measured along the transect by noting the point along the tape where the canopy began and the point at which it ended for each woody species over 1 m tall. Because species overlapped in some cases, the sum of the cover for all species did not necessarily reflect the actual percentage of overstory cover along the tape. The percentage of the tape covered by overstory was also calculated. The height of the tallest vegetation within each continuous stretch of the same species was measured. Woody stem density was determined by using a meter stick to measure one meter outward on the upstream side of the transect. All woody stems within the one meter belt transect were counted and recorded by species into 4 size classes.

Hydrologic variables, including discharge and depth to groundwater as they relate to vegetation, will also be incorporated in the monitoring program. Piezometers were installed within the floodplain near three of the vegetation monitoring sites to provide hydrologic data for correlation analysis. Installation occurred in November 2012 in the Eastside Bypass near Transects 1 and 2, in January 2013 in Reach 4A, and in February 2013 in Reach 2B. A riparian systems model was used to rank riparian habitat condition. This qualitative model (riparian rank) includes spatial and structural diversity of native woody plants, contiguity of dominant vegetation, invasive vegetation, hydrology, topographic complexity, characteristics of flood-prone areas, and biogeochemical processing. These criteria consider the interaction between geology, hydrology, and organic and inorganic inputs to the system.

This page left blank intentionally.

RECLAMATION

Managing Water in the West

Central Valley Steelhead Monitoring Plan for the San Joaquin River Restoration Area

2013 Monitoring Results for National Marine Fisheries Service
Permit 16608



U.S. Department of the Interior
Bureau of Reclamation
Mid-Pacific Region and
Denver Technical Service Center

May 2013

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE May 2013		2. REPORT TYPE Report		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Central Valley Steelhead Monitoring Plan for the San Joaquin River Restoration Area				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Donald E. Portz, Shaun Root, and Charles Hueth				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Bureau of Reclamation, Technical Service Center Fisheries and Wildlife Resources Group PO Box 25007, Denver, CO 80225-0007				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Bureau of Reclamation, Mid-Pacific Region Sacramento, CA 95825				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Available from the National Technical Information Service (NTIS) Operations Division, 5285 Port Royal Road, Springfield, VA 22161					
13. SUPPLEMENTARY NOTE					
14. ABSTRACT Steelhead abundance and distribution in the San Joaquin River (SJR) Basin have substantially decreased, and steelhead are now believed to be extirpated from the Restoration Area. Central Valley (CV) steelhead distinct population segment includes tributaries to the SJR and therefore the presence of CV steelhead must be monitored. Interim flows could attract adult steelhead into the Restoration Area and attracted fish would not have access to appropriate spawning habitat due to a number of impassable barriers. The Bureau of Reclamation implemented a steelhead monitoring and detection plan for the SJR upstream of the Merced River confluence that would, in the event of a capture, document and transport the fish to suitable habitats downstream from the mouth of the Merced River. Electrofishing, fyke traps, and trammel netting collection methods were used for detection of CV steelhead from approximately 3.5 miles upstream of Highway 165 Bridge to the confluence of the Merced River and adjoining sloughs. A total of 27 fish species comprised of 1,248 individuals were captured during Steelhead Monitoring Plan (SMP) activities from January 2013–March 2013. During the SMP period no steelhead were detected in the study area. However, ancillary data that were collected is valuable in that it is the foundational information of fish community assemblages and native fishes for Reach 5 of the San Joaquin River Restoration Program (SJRRP). Six out of 27 fish species captured were native to the SJR, but native fish species only comprised 6.4 percent of total individuals captured. Continued monitoring of adult CV steelhead migration in the Restoration Area is important to provide information regarding the progress of SJRRP while helping to protect fisheries.					
15. SUBJECT TERMS Steelhead, <i>Oncorhynchus mykiss</i> , Steelhead Monitoring Plan, San Joaquin River, San Joaquin River Restoration Program					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 27	19a. NAME OF RESPONSIBLE PERSON Donald E. Portz, Ph.D.
a. REPORT	b. ABSTRACT	a. THIS PAGE			19b. TELEPHONE NUMBER (Include area code) 303-445-2220

Central Valley Steelhead Monitoring Plan for the San Joaquin River Restoration Area

**2013 Monitoring Results for National Marine Fisheries Service
Permit 16608**

by

Donald E. Portz, Shaun Root, and Charles Hueth

Bureau of Reclamation
Denver Technical Service Center
Fisheries and Wildlife Resources Group, 86-68290
PO Box 25007
Denver, CO 80225-0007



**U.S. Department of the Interior
Bureau of Reclamation
Mid-Pacific Region and
Denver Technical Service Center**

May 2013

MISSION STATEMENTS

The U.S. Department of the Interior protects America's natural resources and heritage, honors our cultures and tribal communities, and supplies the energy to power our future.

The mission of the Bureau of Reclamation is to manage, develop, and protect water and related resources in an environmentally and economically sound manner in the interest of the American public.

DISCLAIMER

Mention of commercial products does not constitute endorsement.

TABLE OF CONTENTS

	<i>Page</i>
Abstract.....	v
Introduction.....	1
Central Valley Steelhead.....	3
Study Area	3
Methods	5
Sampling Method 1: Electrofishing.....	6
Sampling Method 2: Fyke Traps with Wing Walls	7
Sampling Method 3: Steelhead Specific Trammel Nets.....	8
Fish Handling and Relocation.....	9
Results.....	10
Electrofishing.....	10
Fyke Traps with Wing Walls	12
Trammel Nets.....	14
Combined Sampling Results.....	15
Discussion.....	17
Acknowledgments	18
References.....	18

Tables

<i>Table</i>	<i>Page</i>
1 Boat electrofishing per unit of time per number of fish captured.....	12
2 Total of fyke netting hours per number of fish captured by location.....	14
3 Trammel netting total (fish per 10,000 net ft/hr)	16
4 Percentage of total fish captures by method	17

Figures

<i>Figure</i>	<i>Page</i>
1 Reaches within the San Joaquin River Restoration Area, San Joaquin Watershed, Central Valley, California.	4
2 Overview of Reach 5 of the San Joaquin River Restoration Area.....	5

Figures (continued)

<i>Figure</i>		<i>Page</i>
3	Reclamation biologist sampling fish with an electrofishing raft.	6
4	Reclamation biologist deploying a fyke net in Mud Slough, Reach 5 of the San Joaquin River Restoration Area.	7
5	Reclamation biologist inspecting a fyke net at Hills Ferry Barrier location near Newman, California.	8
6	Trammel net including an illustration of the pockets that are created by the entangled fish.	9
7	Reclamation biologist retrieving a trammel net on the San Joaquin River, California.	9
8	Fish captured during 2013 Steelhead Monitoring Plan electrofishing sampling.	11
9	Locations of boat electroshocking throughout Reach 5 of the San Joaquin River, California.	11
10	Fish captured during Steelhead Monitoring Plan fyke sampling.	12
11	Fyke net locations throughout Reach 5 of the San Joaquin River Restoration Area, California.	13
12	Fish captured during Steelhead Monitoring Plan using trammel net sampling, 2013.	14
13	Trammel netting locations in Reach 5 of the San Joaquin River, California.	15
14	Total number of fish captured by fish species.	16

ABSTRACT

Steelhead abundance and distribution in the San Joaquin River (SJR) Basin have substantially decreased, and steelhead are now believed to be extirpated from the Restoration Area. Central Valley (CV) steelhead distinct population segment includes tributaries to the SJR and therefore the presence of CV steelhead must be monitored. Interim flows could attract adult steelhead into the Restoration Area and attracted fish would not have access to appropriate spawning habitat due to a number of impassable barriers. The Bureau of Reclamation implemented a steelhead monitoring and detection plan for the SJR upstream of the Merced River confluence that would, in the event of a capture, document and transport the fish to suitable habitats downstream from the mouth of the Merced River. Electrofishing, fyke traps, and trammel netting collection methods were used for detection of CV steelhead from approximately 3.5 miles upstream of Highway 165 Bridge to the confluence of the Merced River and adjoining sloughs. A total of 27 fish species comprised of 1,248 individuals were captured during Steelhead Monitoring Plan (SMP) activities from January 2013–March 2013. During the SMP period no steelhead were detected in the study area. However, ancillary data that were collected is valuable in that it is the foundational information of fish community assemblages and native fishes for Reach 5 of the San Joaquin River Restoration Program (SJRRP). Six out of 27 fish species captured were native to the SJR, but native fish species only comprised 6.4 percent of total individuals captured. Continued monitoring of adult CV steelhead migration in the Restoration Area is important to provide information regarding the progress of SJRRP while helping to protect fisheries.

INTRODUCTION

In 1988, a coalition of environmental groups, led by the Natural Resources Defense Council (NRDC), filed a lawsuit challenging the renewal of long-term water service contracts between the United States and the Central Valley Project Friant Division Long-Term Contractors. After more than 18 years of litigation of this lawsuit, known as *NRDC, et al. v Kirk Rodgers, et al.*, a settlement was reached. On September 13, 2006, the Settling Parties, including NRDC, Friant Water Users Authority, and the U.S. Departments of the Interior and Commerce, agreed on the terms and conditions of the settlement, which was subsequently approved by the U.S. Eastern District Court of California on October 23, 2006. The Settlement establishes two primary goals: (1) Restoration Goal – To restore and maintain fish populations in “good condition” in the mainstem San Joaquin River below Friant Dam to the confluence of the Merced River, including naturally reproducing and self-sustaining populations of salmon and other fish, and (2) Water Management Goal – To reduce or avoid adverse water supply impacts on all of the Friant Division long-term contractors that may result from the Interim Flows and Restoration Flows provided for in the Settlement. These goals will require developing a fisheries management plan that implements an adaptive management approach that includes professional environmental review, review of structural modifications and designs, and technical support to provide the best quality data to define problems, prioritize actions, and increase the confidence in future decisions.

Potential routes to spawning habitats for migratory fish such as the Central Valley (CV) steelhead (*Oncorhynchus mykiss*) are believed to have been historically unhindered in the San Joaquin River (SJR) before completion of the Friant Dam. Although little detailed information on steelhead distribution and abundance is available (Lindley *et al.* 2006, McEwan 2001), they are mostly distributed higher in watersheds with large river systems than Chinook salmon (*Oncorhynchus tshawytscha*; Voight and Gale 1998, as cited in McEwan 2001). Therefore, steelhead may have spawned at least as far upstream as the natural barrier located at the present-day site of Mammoth Pool and the upper reaches of SJR tributaries. Modeling of potential steelhead habitat by Lindley *et al.* (2006) suggests that a portion of the upper SJR basin historically supported an independent steelhead population. However, much of the habitat downstream from this population’s modeled distribution may have been unsuitable for rearing because of high summer water temperatures. Lindley *et al.* (2006) concluded that suitable steelhead habitat existed historically in all major SJR tributaries, although to a lesser degree than in stream systems in the Cascades, Coast Range, and Northern Sierra Nevada. Additionally, steelhead are historically documented in the Tuolumne and Kings River systems (McEwan 2001).

Steelhead abundance and distribution in the SJR basin have substantially decreased (McEwan 2001), and steelhead have been extirpated from the Restoration Area after the construction of Friant Dam. Based on their review

of factors contributing to steelhead declines in the Central Valley, McEwan and Jackson (1996) concluded that basin-wide population declines were related to water development and flow management that resulted in habitat loss. Dams have blocked access to historical spawning and rearing habitat upstream, thus forcing steelhead to spawn and rear in the lower portion of the rivers where water temperatures are often high enough to be lethal (Yoshiyama *et al.* 1996, McEwan 2001, Lindley *et al.* 2006). However, steelhead continue to persist in low numbers in the Stanislaus, Tuolumne, and Merced River systems (McEwan 2001, Zimmerman *et al.* 2008). CV steelhead distinct population segment (DPS); smallest division of taxonomic species protected under the U.S. Endangered Species Act; 61 FR 4722) includes tributaries to the SJR that drain the western slopes of the Sierra Nevada Mountains (*i.e.*, Mokelumne, Calaveras, Stanislaus, Tuolumne, Merced, Chowchilla, Fresno, upper San Joaquin, Kings, Kaweah, and Kern Rivers, and Caliente Creek; NMFS 2009).

Monitoring of CV steelhead populations in the SJR and its tributaries is especially challenging due to extremely low abundance of fish. CV steelhead populations are depressed to the point where monitoring opportunities are limited because sample sizes are too low to use statistical analyses (Eilers *et al.* 2010), and depressed to the point that even determination of presence is difficult. According to Eilers *et al.* (2010), CV steelhead are currently extirpated from all waters upstream of the Merced-San Joaquin River confluence. However, annual spring interim flows occurring from February 1–June 1 could attract adult steelhead into the Restoration Area. Attracted steelhead would not have access to appropriate spawning habitat due to a number of impassable barriers. Therefore, the Bureau of Reclamation (Reclamation) implemented a steelhead monitoring and detection plan (SMP) for the SJR upstream of the Merced River confluence that would, in the event of a capture, document and transport the fish to suitable habitats downstream from the mouth of the Merced River.

Annual fall interim flows occurring from October 1–December 1 could also attract adult steelhead into the Restoration Area if the interim flows are higher than the flows in the SJR tributaries. However, during fall interim flows, the Hills Ferry Barrier (HFB) is in place just upstream of the confluence with the Merced River and ongoing fish monitoring occurs at HFB. Steelhead that reach the HFB could be detected and potentially trapped or deterred from upstream migration. In the fall of 2010 and 2011, a trap was operated by the Reclamation, Denver Technical Service Center's fisheries biologist to assess the barrier's effectiveness. Some fall-run Chinook salmon were able to pass the barrier during the 2010 and 2011 interim flow period, so the effectiveness of HFB is in question (Portz *et al.* 2011). No steelhead were detected, however bar spacing on the trap could allow steelhead that are smaller and slimmer than salmon to escape.

Central Valley Steelhead

Steelhead are the anadromous form (*i.e.*, returning from sea to the river in order to spawn) of *Oncorhynchus mykiss*. This anadromous fish is listed under the federal Endangered Species Act as threatened species by the National Marine Fisheries Service (NMFS; NMFS 1998). Critical habitat was designated for CV steelhead in 2005 (70 FR 52488). The designated critical habitat for CV steelhead in the San Joaquin River Basin include the Mokelumne, Calaveras, Stanislaus, Tuolumne, and Merced Rivers. On August 15, 2011, NMFS completed the 5-year status review of CV steelhead and recommended that CV steelhead DPS remain classified as a threatened species. Currently, CV steelhead DPS and critical habitat extends from the San Joaquin River to the confluence with the Merced River (NMFS 2011).

Steelhead in the California Central Valley are divided into two types: summer-run and winter-run. Summer-run steelhead are river-maturing fish species that require coldwater pools between 55°F and 70°F for holding and staging. According to Lindley *et al.* (2006), summer-run steelhead have been extirpated because suitable summer holding habitats are located above impassable dams. Therefore, ocean maturing winter-run steelhead is the only type found in the Central Valley (Moyle 2002).

Two to three year-old CV steelhead generally migrate to freshwater (Reynolds 1993), and occurrence of adults in the San Joaquin River range between July and March of the following year, but peaks between the months of December and January (CDFG 2007) when small streams and tributaries are cool and well-oxygenated (Williams 2006). Unlike other salmonids which can only spawn once before death, a percentage of steelhead population (17.2 percent) in California streams can return to the ocean and migrate back upstream to spawn again in subsequent years (Shapolov and Taft 1954).

STUDY AREA

The Restoration Area for the San Joaquin River Restoration Program (SJRRP) includes the SJR between Friant Dam and its confluence with the Merced River (Figure 1). Steelhead monitoring activities were proposed as the area of the SJR below Sack Dam to the confluence with the Merced River and adjoining sloughs. For this study, Sack Dam was considered the furthest upstream extent for CV steelhead migration because it is impassable except during flood conditions. During the winter of 2013 monitoring was confined to Reach 5 and the lower extent of 4B2 of the San Joaquin River Restoration Area (Figure 2) because SJRRP made no flow releases below Sack Dam, drying major portions of the Reach 4A channel. A total of approximately 18 river miles along the San Joaquin River were monitored as well as slough tributaries (totaling

approximately 19.4 river miles) for a total of approximately 37.4 river miles monitored (Figure 2).



Figure 1.—Reaches within the San Joaquin River Restoration Area, San Joaquin Watershed, Central Valley, California.




Figure 2.—Overview of Reach 5 of the San Joaquin River Restoration Area, defined as Salt Slough Confluence, Grasslands State Park, Mud Slough Confluence, Newman Wasteway, Merced River Confluence. Tributaries pertaining to steelhead monitoring are highlighted in red.

METHODS

Migrating adult steelhead are difficult to monitor with commonly used salmonid monitoring techniques (*e.g.*, carcass surveys, snorkel surveys, redd counts) due to their unique life-history traits. Steelhead, unlike salmon, may not die after spawning. Therefore, carcasses may not be available for a mark-recapture survey. In addition, steelhead migrate and spawn during the late-fall, winter, and spring months when rivers have periods of pulse flows (*e.g.*, Vernalis Adaptive Management Plan or VAMP), high flows (*e.g.*, flood releases), and turbid water conditions. Three sampling methods were implemented for this CV steelhead monitoring plan:

Sampling Method 1: Electrofishing

Electrofishing is a common method used in monitoring steelhead population (e.g., Mill and Deer creeks, and Feather, American, Mokelumne, Stanislaus, and Merced rivers). One potential drawback for using electrofishing in rivers involves the difficulty in obtaining permits due to the possibility of injuring fish in anadromous salmonid waters (Eilers 2008). However, electrofishing effectiveness and safety have improved over time (Bonar *et al.* 2009). Design specifications to reduce injury to fish, and a comprehensive review of electrofishing literature can be found in Snyder (2003).

Sampling was done monthly from January–March 2013. Capture of resident fish multiple times was to be anticipated, thus monthly sampling was important to ensure fish recovery from sampling and handling stress between captures. Electrofishing methods followed the NMFS guidelines for sampling waters with anadromous fish. However, stated guidelines were for backpack electrofishing, but SMP researchers were not precluded from using boat electrofishing. NMFS were given substantial proof that the proposed techniques and equipment were necessary for the study and that listed species were safeguarded (NMFS 2000) and state scientific collecting permits were obtained for these sampling techniques.

A Reclamation raft electroshocker was used on January 14-15, 2013, February 12-13, 2013, and March 6-7, 2013 with a crew of Reclamation staff (Figure 3). The settings were set at low 50-500 volts with an average power of 55 percent depending on water conductivity and a direct current at 30 pulses per second to ensure that electrical injury to fish was minimized.



Figure 3.—Reclamation biologist sampling fish with an electrofishing raft.

Sampling Method 2: Fyke Traps with Wing Walls

Fyke nets with wing walls and traps (Figures 4 and 5) were used to sample upstream migrating CV steelhead. Fyke nets have long been used to capture migrating fish to monitor their yearly changes in abundance. The nets were constructed of 2.38-cm square #252 knotless nylon netting formed over 5 consecutive 1.22-m hoops and a 1.2 x 1.2 m welded conduit frame entrance. The traps contained 2 throats with a 25 cm diameter opening. Wings walls were 1.22 m deep and long enough to span the river (max wing length 30.5 m) with small floats spaced every 61 cm on top and a lead line on bottom. Nets were held in place with anchored t-posts and were deployed in sampling locations (*i.e.*, upstream of the confluence of the Merced River, the mouths of Mud Slough, Newman Wasteway, Van Clief Road, and within Salt Slough). This technique was implemented once the Hills Ferry Barrier was removed in mid-December 2012. Fyke nets were deployed at sampling locations until March 15, 2013. Marking buoys and flashing amber lights were affixed for safety and to alert boaters of the net's presence. The traps were checked daily so the likelihood of fish being physically injured was low. In the event of a steelhead capture, the fish would have been measured, weighed, transported, and released downstream from the mouth of the Merced River.



Figure 4.—Reclamation biologist deploying a fyke net in Mud Slough, Reach 5 of the San Joaquin River Restoration Area.



Figure 5.—Reclamation biologist inspecting a fyke net at Hills Ferry Barrier location near Newman, California.

Fyke nets were used in lieu of wire fyke traps for several reasons: fyke nets were relatively inexpensive and easy to install, not a boat passage impediment (can be pushed down in the water column for boat passage), easily replaced if damaged, easily transported, and no permitting was required to transport. Although California Department of Fish and Wildlife (CDFW) owns a wire fyke trap capable of catching fish in high flows, it was not used because it would have required a crane to remove the trap out of water under increased hydraulic pressure and in the event that the trap became silted.

Sampling Method 3: Steelhead Specific Trammel Nets

Trammel nets (Figure 6) are most common as stationary gear to block off channels with low velocities or no flows. However, they can also be used to drift in short durations (*e.g.*, 20 min) in moving water currents. For this study, short duration drifting of trammel nets were deployed as well as stationary sets. Short fishing durations prevent fish from being severely entangled and lessen the chance of harm. Trammel nets were advantageous and relatively efficient in turbid waters. The nets consisted of three parallel vertical layers of netting; the inner net had a very small mesh size, while the outer nets had mesh size large enough for fish to pass. The larger and smaller mesh size nets form a pocket when fish try to swim through (Figure 6). Similar to seine nets, trammel nets were equipped with




Figure 6.—Trammel net including an illustration of the pockets that are created by the entangled fish.



Figure 7.—Reclamation biologist retrieving a trammel net on the San Joaquin River, California.

floats attached to the head rope and lead weights along the ground rope. Colored floats were attached to the head rope so boaters and other recreationists could be alerted to the nets and avoid entangling themselves, their boats, and/or their fishing gear. To ensure safety of steelhead, fisheries biologists tending the nets followed at a close distance to observe, reduce risk of entanglement, and retrieve nets in short time intervals. Sampling time depended on the number of fish and by-catch caught at each location. Trammel net sampling was used during adult steelhead migration January–March, 2013 on the SJR. It should also be noted that although the data is broken into two seemingly different projects *i.e.*, Steelhead Monitoring Plan and Inventory and Monitoring, the methods and implementation are the same; therefore the data is within the scope and permitting of the Steelhead Monitoring Plan.

Fish Handling and Relocation

In the event that CV steelhead were captured during monitoring activities, fish

would have been subjected to standard handling and transporting procedures. Captured steelhead would be recorded, measured (*i.e.*, fork length and total length), sexed (if possible), sampled for scales and tissues, and checked for injuries and presence of tags. Additionally, fish would have been Floy tagged with a unique identification number to document any recaptures that may occur in the study area. Captured steelhead were to be transported downstream from the mouth of the Merced River in a 550-L transport tank. The transport tank would have been immediately filled with river water prior to transport using a portable screened water pump. Steelhead were to be transferred from the river to the transport tank using a water-to-water transfer to help minimize stress and loss of slime. Oxygen gas was to be supplied using a compressed oxygen gas cylinder and micro-bubble diffusers to maintain dissolved oxygen levels near saturation. Sodium chloride would have been added to the transport water to decrease the cellular-holding water ionic gradient as a means to minimize stress. The truck was to be stopped after 30 minutes of transportation and each hour thereafter for visual inspection of the fish and transport equipment. Captured CV steelhead were to be acclimated to receiving water temperature and water quality at the predetermined release location.

RESULTS

No CV steelhead were observed during this monitoring period in the SJR Restoration Area.

Electrofishing

Raft electrofishing yielded 23 fish species (Figure 8), representing 54 percent ($n=673$) of the cumulative total ($n=1248$) of all individuals captured across the three methods (see Table 4). Non-native fishes made up 95 percent ($n=637$) of individuals captured using this method. Native species comprised only 5 percent ($n=36$) of fish captured with this method, and 3 percent of the cumulative total. Electrofishing was the most successful methodology used, based on sheer numbers of individuals captured, as well as the diversity of individuals captured. No CV Steelhead were captured with this sampling method.

Selected locations within the study area were raft electrofished during the sampling season (Figure 9). The total generator shocking time was 21,211 sec (avg. 542.3 sec/location), and a total of 673 fish captured. Therefore, the catch per unit effort (CPUE) was 1.9 fish per minute (Table 1).




Figure 8.—Fish captured during 2013 Steelhead Monitoring Plan electrofishing sampling.




Figure 9.—Locations of boat electroshocking throughout Reach 5 of the San Joaquin River, California. Markers represent approximate start locations as some of the markers represent multiple shocking locations at this resolution.

Table 1.—Boat electrofishing per unit of time per number of fish captured

Boat Electrofishing Total (n=673)				
Electrofishing	Seconds	Minutes	Fish Captured	CPUE per Minute
Steelhead Monitoring	14,342	239.03	406	1.699
Inventory and Monitoring	6,869	114.48	267	2.332
Total	21,211	353.51	673	1.904

Fyke Traps with Wing Walls

There were 18 fish species caught totaling 393 individuals (Figure 10). Fyke netting represented 31 percent of the cumulative total (n=1248) of all individuals captured. Non-native fishes were 98 percent (n=384) of individuals captured using this method, and 31 percent of the cumulative total. Native species comprised only 2 percent (n=9) of fish captured with this method, and 0.7 percent of the cumulative total. No CV Steelhead were captured using this method.




Figure 10.—Fish captured during Steelhead Monitoring Plan fyke sampling.

Five fyke nets were deployed throughout the study area from approximately 2.8 river miles upstream of the Highway 165 Bridge to near the SJR confluence with the Merced River. Nets were set at: (1) Hills Ferry Barrier location, (2) Mud Slough, (3) Newman Wasteway, (4) Salt Slough, (5) and Van Clief (Figure 11).




Figure 11.—Fyke net locations throughout Reach 5 of the San Joaquin River Restoration Area, California.

Fishing hours differed between nets due to net damage or dangerous flows resulting in safety concerns, however when in the water, nets fished for approximately 24 hours each day from January 17–24, 2013, February 4–14, 2013, and March 4–14, 2013.

The five fyke locations fished for a combined total of 116.6 days during the sampling period. However, beavers (*Castor canadensis*) chewed through the fyke net traps on 13 different occasions. For this reason, 13.5 days will not be incorporated to the total days of fyke net deployments. Fishing hours for fyke nets exclusive of beaver damage total 103 days (2,473.3 hours). The CPUEs for each trapping location are provided in Table 2. The nets had a cumulative average of 3.8 fish per trap per day (Table 2).


Table 2.—Total of fyke netting hours per number of fish captured by location

Fyke Netting Exclusive of Beaver Damage				
Location	Hours Fished	Fish Captured	Days Fished	Fish per day
Hills Ferry	549.04	8	22.88	0.35
Mud Slough	251.15	20	10.47	1.91
Newman Wasteway	606.55	124	25.27	4.91
Salt Slough	490.2	26	20.43	1.27
Van Clief	576.33	215	24.01	8.95
Fyke Net total	2473.27	393	103.05	3.82

Trammel Nets

Trammel netting accounted for a total of 18 species (Figure 12), representing 15 percent (n=182) of the cumulative total (n=1248) of all individuals captured across the three methods (see Table 4). Non-native fishes captured made up 84 percent (n=152) of individuals captured using this method, and 12 percent of the cumulative total. Native species comprised only 16 percent (n=30) of fish captured with this method, and 2 percent of the cumulative total. No CV Steelhead were captured using this method.

Figure 12.—Fish captured during Steelhead Monitoring Plan using trammel net sampling,



2013.

Trammel nets were deployed or drifted opportunistically throughout Reach 5 of the Restoration Area from January 17–24, 2013, February 4–14, 2013, and March 4–14, 2013 (Figure 13). These nets were measured in feet that varied in sizes: 3 × 40, 3 × 50, 4 × 50, 4 × 75, 5 × 50, 5 × 100, and 6 × 50. A total of 59 net deployments with varying sizes occurred during sampling period. The total deployed square footage was 29,535 ft² with 1967.7 total hours of fishing time. Therefore, the cumulative CPUE is 0.03 fish per 10,000 ft²/hr (Table 3).




Figure 13.—Trammel netting locations in Reach 5 of the San Joaquin River, California. Markers represent approximate locations and may represent multiple nets or nets in close proximity at this resolution.

Combined Sampling Results

A total of 27 fish species comprised of 1,248 individuals were captured during the entire sampling period (Figure 14). Non-native fishes were 94 percent (n=1173) of all fish captured with the common carp (20.9 percent), bluegill (16.5 percent),

Table 3.—Trammel netting total (fish per 10,000 net ft/hr)

Trammel Netting Total					
	Hours Fished	Net Size (ft ²)	Fish Captured	Fish per 10,000 ft ²	Fish per 10,000 net ft ² /hr
Steelhead Monitoring	709.52	10216	120	117.46	0.166
Inventory and Monitoring	1258.14	19319	62	32.09	0.026
Total	1967.66	29535	182	61.62	0.031




Figure 14.—Total number of fish captured by fish species.

and goldfish (11.5 percent) comprising almost 50 percent (n=573) of non-native fish captured (n=1173), and 46 percent (n=1248) of all fish captured (Table 4). Native species were only 6 percent (n=75) of individuals captured during the sampling period. Native species largely consisted of Sacramento blackfish (40 percent), Sacramento sucker (36 percent), and Sacramento splittail

Table 4.—Percentage of total fish captures by method

Method	Percentage of Non-Native Fish	Percentage of Native Fish	Percentage of Total Fish Captured
Boat Electroshocking	54% (n=637)	48% (n=36)	54% (n=673)
Fyke Net	33% (n=384)	12% (n=9)	31% (n=393)
Trammel Net	13% (n=152)	40% (n=30)	15% (n=182)
Total	94% (n=1173)	6% (n=75)	100% (n=1248)

(13 percent) that summed a cumulative total of 93 percent of native fish captures, and 5.6 percent (n=70) of all fish (see Table 4). A male Chinook salmon was captured at Mud Slough using a fyke net January 20, 2013.

DISCUSSION

Historically, the SJR Restoration Area was a potential migratory pathway for CV steelhead to reach their spawning grounds. However, little detailed information on their distribution and abundance is available for these river reaches. The upper SJR basin may have historically supported a steelhead population but much of the downstream habitat is unsuitable for rearing because of high summer water temperatures (Lindley *et al.* 2006). Suitable steelhead habitat existed historically in all major SJR tributaries and there is potential for this fish to return under interim flows. Annual fall interim flows occurring from October 1 to December 1 and spring interim flows occurring from February 1 to June 1 could attract adult steelhead into the restoration area. Attracted steelhead would not have access to appropriate spawning habitat due to a number of impassable barriers. However this is thought to be relatively unlikely because CV steelhead are currently extirpated from all waters upstream of the Merced-San Joaquin River confluence (Eilers *et al.* 2010).

Monitoring of CV steelhead populations in the SJR and its tributaries is especially challenging due to extremely low abundance or absence of fish. During the SMP period no steelhead were detected in the study area. However, ancillary data that were collected is valuable in that it is the foundational baseline information of fish community assemblages and native fishes for Reach 5 of the SJRRP. Six out of 27 fish species captured were native to the SJR, though populations of these native fish species were much less represented compared to non-native. Only 6 percent of total individuals captured were native to California waters.

Lastly, the steelhead monitoring plan is an important study for the SJRRP to

ensure its commitment to restore and maintain fish populations within the Restoration Area. Although no CV steelhead were detected or captured during this sampling period, the continued monitoring of adult CV steelhead migration in the Restoration Area is important to provide information regarding the progress of the Restoration Program. Monitoring population abundance trends, rare and native species occurrences, and fish community assemblages will provide a biological indication of SJRRP's success.

ACKNOWLEDGMENTS

We would like to thank Alicia Forsythe, Michelle Banonis, and Erin Rice for their program level support and guidance. We would also like to thank Matt Bigelow and staff of the California Department of Fish and Wildlife, Leslie Mirise of the National Marine Fisheries Service, and Eric Best, Durel Carstensen, Jarod Hutcherson, Juddson Sechrist, Zak Sutphin of the Bureau of Reclamation Technical Service Center for their assistance with monitoring and data collection. We would also like to acknowledge Erin Rice of the San Joaquin River Restoration Program Office for his assistance with permitting. Funding for this monitoring was provided by the Bureau of Reclamation Mid-Pacific Region's San Joaquin River Restoration Program.

REFERENCES

- Bonar, S.A., W.A. Hubert, and D.W. Willis, eds., 2009. Standard Methods for Sampling North American Freshwater Fishes. American Fisheries Society. Bethesda, Maryland. 335 p.
- California Department of Fish and Game (CDFG). 2007. California Steelhead Fishing Report-Restoration Card. A Report to the Legislature. July 2007.
- Eilers, C.D. 2008. Review of present steelhead monitoring programs in the California Central Valley. Sacramento, California. Report to California Department of Fish and Game Contract # P0685619.
- Eilers, C.D., J. Bergman, and R. Nelson. 2010. A Comprehensive Monitoring Plan for Steelhead in the California Central Valley. The Resources Agency: Department of Fish and Game: Fisheries Branch Administrative Report Number: 2010-2.

- Lindley S.T., R.S. Schick, A. Agrawal, M. Goslin, T. Pearson, E. Mora, J.J. Anderson, B. May, S. Greene, C. Hanson, A. Low, D. McEwan, R.B. MacFarlane, C. Swanson, and J.G. Williams. 2006. Historical population structure of Central Valley steelhead and its alteration by dams. *San Francisco Estuary and Watershed Science* 4, Issue 1, Article 2.
- McEwan, D.R. 2001. Central Valley Steelhead IN: Contributions to the biology of central valley salmonids. *Fish Bulletin* 179.
- McEwan, D. and T.A. Jackson. 1996. Steelhead restoration and management plan for California. California Department of Fish and Game, Sacramento. 234 pp.
- Moyle, P.B. 2002. *Inland Fishes of California*. University of California Press, Berkeley.
- National Marine Fisheries Service (NMFS). 1998. Endangered and Threatened Species: Proposed endangered status for two Chinook salmon ESUs and proposed threatened status for five Chinook salmon ESUs; Proposed redefinition, threatened status, threatened status, and revision of critical habitat for one Chinook salmon ESU; Proposed designation of Chinook salmon critical habitat in California, Oregon, Washington, Idaho. March 9, 1998. *Federal Register* 63 (45):11482–11520.
- National Marine Fisheries Service (NMFS). 2000. Guidelines for Electrofishing Waters Containing Salmonids Listed Under the Endangered Species Act.
- National Marine Fisheries Service (NMFS). 2011. Endangered and Threatened Species; Take of Anadromous Fish. *Federal Register* 76:77490–77491.
- National Marine Fisheries Service (NMFS). 2009. Public Draft Recovery Plan for the Evolutionarily Significant Units of Sacramento River Winter-run Chinook Salmon and Central Valley Spring-run Chinook Salmon and the Distinct Population Segment of Central Valley Steelhead. Sacramento Protected Resources Division. October 2009.
- Portz D.E., E. Best, and C. Svoboda. 2011. Evaluation of Hills Ferry Barrier effectiveness at restricting Chinook salmon passage on the San Joaquin River. U.S. Department of Interior, Bureau of Reclamation, Technical Service Center.
- Reynolds, F.L., T.J. Mills, R. Benthin, and A. Low. 1993. Restoring Central Valley streams: a plan for action. California Department of Fish and Game, Inland Fisheries Division, Sacramento.

- Shapovalov, L. and A.C. Taft. 1954. The life histories of the steelhead rainbow trout (*Salmo gairdneri gairdneri*) and silver salmon (*Oncorhynchus kisutch*) with special reference to Waddell Creek, California, and recommendations regarding their management. California Department of Fish and Game, Fish Bulletin 98. 375 p.
- Snyder, D.E. 2003. Electrofishing and its harmful effects on fish. US Geological Survey, Information and Technology Report USGS/BRD/IRT-2003-0002, Denver, Colorado.
- Voight, H.N. and Gale D.B. 1998. Distribution of fish species in tributaries of the lower Klamath River: an interim report, FY 1996. Technical Report No. 3. Yurok Tribal Fisheries Program, Habitat Assessment and Biological Monitoring Division. 71 p.
- Williams, J.G. 2006. Central Valley salmon: a perspective on Chinook and steelhead in the Central Valley of California. San Francisco Estuary and Watershed Science 4(3): Article 2. 416 p.
- Yoshiyama, R.M., E.R. Gerstung, F.W. Fisher, and P.B. Moyle. 1996. Historical and present distribution of Chinook salmon in the Central Valley drainage of California, Sierra Nevada Ecosystem Project: final report to congress, Volume III: Assessments, commissioned reports, and background information, University of California, Center for Water and Wildlife Resources, Davis, CA, pp. 309–362.
- Zimmerman, C.E., G.W. Edwards, and K. Perry. 2008. Maternal origin and migratory history of *Oncorhynchus mykiss* captured in rivers of the Central Valley, California. Final Report. Prepared for California Department of Fish and Game, Contract P03853006. March 2008.
[http://www.delta.dfg.ca.gov/AFRP/documents/Zimmerman_et_al.pdf]